

Ampelomyces quisqualis
(*Cicinnobolus cesatii*)
on Queensland Erysiphaceae

by
B. G. CLARE

University of Queensland Papers

Department of Botany

Volume IV

Number 11

THE UNIVERSITY OF QUEENSLAND PRESS

St. Lucia

10 June 1964

301

· U66

v. 4 no. 11

1

Fryer

WHOLLY SET UP AND PRINTED IN AUSTRALIA BY
WATSON FERGUSON AND COMPANY, BRISBANE, QUEENSLAND

1964

REGISTERED IN AUSTRALIA FOR TRANSMISSION BY POST AS A BOOK

AMPELOMYCES QUISQUALIS (CICINNOBOLUS CESATII) ON QUEENSLAND ERYSIPHACEAE

Ampelomyces quisqualis Ces., *Bot. Zeit.* (1852) **10**: 301.

Cinchnobolus cesatii de Bary, *Abh. senckenb. naturf. Ges.* (1870) **7**: 431 (Beitrage zur Morphologie und Physiologie der Pilze 111: 71), pls. 11, 12, figs. 5-13.

Figures 1 and 2

Hyphae of the hyper-parasite were hyaline and septate; they were present within the hyphae, conidiophores, and conidia of infected mildews. Pycnidia were light brown in transmitted light and varied in shape from sub-globose to pyriform. Pycnidial shape appeared to be determined by the shape of the structure within which they developed; in the *Quercus* mildew they occurred within terminal conidia and were sub-globose, whereas in the other mildews they developed within conidiophores and were ellipsoid or pyriform. Pycnidia contained one-celled, hyaline, guttulate conidia which were ovoid, clavate, cylindrical, or curved. Dimensions are given in Table 1.

TABLE 1
DIMENSIONS OF PYCNIDIA AND CONIDIA IN *Ampelomyces* SPECIMENS

HOST	PYCNIDIA (microns)		CONIDIA (microns)	
	Length	Width	Length	Width
<i>Oidium</i> sp. (<i>Erysiphe cichoracearum</i>) on <i>Cestrum parqui</i>	60-105	30-53	5.5-8.5	1.5-3.0
<i>Oidium</i> sp. (<i>Sphaerotheca fuliginea</i>) on <i>Euphorbia hirta</i>	48-90	30-45	4.0-6.5	2.0-3.5
<i>Oidium</i> sp. (<i>Microsphaera alphitoides</i>) on <i>Quercus pubescens</i>	33-50	30-43	5.5-7.0	2.0-3.0
<i>Oidium</i> sp. (<i>S. fuliginea</i>) on <i>Verbena hybrida</i>	33-66	23-40	4.0-5.5	2.0-4.0
<i>Oidium</i> sp. (<i>E. cichoracearum</i>) on <i>Zinnia elegans</i>	47-83	23-41	4.0-6.5	2.0-4.0




FIG. 1.—*Ampelomyces quisqualis* on *Zinnia elegans* powdery mildew. A. Pycnidia. B. Conidia.



FIG. 2.—*Ampelomyces quisqualis* on *Z. elegans* powdery mildew. Pycnidia (x 600).

Specimens: BRIU* 2041, on *Oidium* sp. (*Erysiphe cichoracearum* DC.), on *Cestrum parqui* L'Hérit., Brisbane, July 1957; BRIU 1951, on *Oidium* sp. (*Sphaerotheca fuliginea* (Schlecht.) Poll.), on *Euphorbia hirta* L., Brisbane, May 1957; BRIU 2056, on *Oidium* sp. (*Microsphaera polonica* Siem.), on *Hydrangea macrophylla* (Thunb.) Ser., R. F. Langdon, Brisbane, July 1959; BRIU 2044, on *Oidium* sp. (*Microsphaera alphitoides* Griff. & Maubl.), on *Quercus pubescens* Willd., R. F. Langdon, Brisbane, October 1959; BRIU 2051 on *Oidium* sp. (*S. fuliginea*), on *Verbena hybrida* Voss., Brisbane, May 1958; BRIU 2045, on *Oidium* sp. (*E. cichoracearum*), on *Zinnia elegans* Jacq., Brisbane, April 1957.

Pycnidia in the mildew on *Hydrangea* were immature and contained no conidia.

In de Bary's description of the species, pycnidial dimensions are 25–35 x 9–15 μ , and conidial dimensions 2.5–3.0 x 1 μ (Saccardo, 1884). However Diedecke claims that pycnidia are 25–60 x 10–15 μ , and conidia 5–10 x 2–4 μ (Grove, 1935).

De Bary was the first to realise that the species was hyper-parasitic and not part of the mildew life-cycle and he named the species *Cicinnobolus cesatii*. This name is the one generally used for the species, but Rogers (1959) has pointed out that it is illegitimate and that *Ampelomyces quisqualis* Ces. should be used instead.

Many species of the genus *Cicinnobolus* have been described. All are parasitic on mildews and all have allegedly distinctive host preferences and morphology, but Blumer (1933) investigated fourteen of these species experimentally and concluded that they showed no distinctive host preferences and were indistinguishable. It would seem that unnecessary synonymy could be avoided if the genus were regarded as being monotypic until further experimental work, such as that of Emmons (1930) and Blumer (1933), has been carried out.

The Queensland specimens are considered to be *Ampelomyces quisqualis* Ces. This is the first record of the species in Queensland on the hosts listed above.

REFERENCES

- BLUMER, S. (1933). Die Erysiphaceen Mitteleuropas mit besonderer Berücksichtigung der Schweiz. *Beitr. Kryptogamenfl. Schweiz* **7**: 1-482.
- EMMONS, C. W. (1930). *Cicinnobolus cesatii*, a study in host-parasite relationships. *Bull. Torrey bot. Cl.* **57**: 421-41.
- GROVE, W. B. (1935). *British Stem and Leaf Fungi*, 1st ed., vol. 1, pp. 150-151. Cambridge: C.U.P.
- SACCARDO, P. A. (1884). *Syll. fung.* **3**: 216.
- ROGERS, D. P. (1959). On *Cicinnobolus*. *Mycologia* **51**: 96-98.

*BRIU=Herbarium, Department of Botany, University of Queensland.